

PRINCIPAL BUNDLES ADMITTING A HOLOMORPHIC CONNECTION

INDRANIL BISWAS

1. INTRODUCTION

Let E be a holomorphic vector bundle over a compact Kähler manifold M , such that E admits a holomorphic connection compatible with its holomorphic structure. The following question is perhaps due to S. Murakami [M1], [M2], [M3] : does there exists a flat connection on E compatible with the holomorphic structure ?

The same question can be posed for a more general principal G bundle where G is a complex Lie group. Murakami constructed examples of torus bundles over torus which admit holomorphic connection but do not admit compatible flat connection; thus providing a negative answer to the general question.

Let M be a compact connected Kähler manifold of complex dimension d , and let ω be a Kähler form on M . The degree of a torsion-free $\mathcal{O}_M$ -coherent sheaf F is defined as [Ko, Ch. V, (7.1)]

$$(1.1) \quad \deg F := \int_M c_1(F) \wedge \omega^{d-1}$$

Consider the Harder-Narasimhan filtration of the holomorphic tangent bundle of M [Ko, page 174, Ch. V, Theorem 7.15] :

$$(1.2) \quad 0 = \mathcal{E}_0 \subset \mathcal{E}_1 \subset \mathcal{E}_2 \subset \mathcal{E}_3 \subset \dots \subset \mathcal{E}_{l-1} \subset \mathcal{E}_l = T$$

Let G be a connected affine algebraic reductive group over $\mathbb{C}$. Let P be a holomorphic principal G bundle on M (*i.e.* the transition functions of P are holomorphic). Assume that P admits a holomorphic connection D compatible with the holomorphic structure. This means the following : D is a holomorphic 1-form on P with values in the Lie algebra, $\mathfrak{g}$, of G , such that D is invariant for the action of G on P , and, when restricted to the fibers of P , this form coincides with the holomorphic Maurer-Cartan form. Using the natural identification of the holomorphic tangent space of P with its real tangent space, the holomorphic connection D gives a G connection on P .

A flat G connection on P is called compatible with respect to the holomorphic structure if the $(1,0)$ component of the connection form (which is a $\mathfrak{g}$ valued 1-form on P) is actually a holomorphic 1-form. This is equivalent to the following : M can be covered by open set $\{U_i\}$ such that over each U_i the principal bundle P admits a holomorphic trivialization which is also constant (with respect to the connection). Clearly the $(1,0)$ component of such a flat connection compatible with the holomorphic structure gives a holomorphic connection compatible with the holomorphic structure.

Supported by the Académie des Sciences.

Our aim here is to prove the following theorem [Theorem 3.1] :

Theorem A. *If the Kähler manifold M satisfies the condition that $\deg(T/\mathcal{E}_{l-1}) \geq 0$, then a holomorphic principal G -bundle P on M admitting a compatible holomorphic connection is semistable. Moreover, if $\deg(T/\mathcal{E}_{l-1}) > 0$, then such a bundle P actually admits a compatible flat G -connection.*

So, in particular, if T is semistable with $\deg T > 0$, then a G -bundle P on M with a holomorphic connection admits a compatible flat connection. In the case where $G = GL(n, \mathbb{C})$, the above result was proved in [Bi].

The author is grateful to D. Akhiezer for some very useful discussions.

2. PRELIMINARIES

We continue with the notation of Section 1.

Let D be a holomorphic connection on the holomorphic principal G bundle P compatible with its holomorphic structure.

Let $Ad(P)$ denote the vector bundle on M associated to P for the adjoint action of G on its Lie algebra $\mathfrak{g}$. The holomorphic structure of P will induce a holomorphic structure on the vector bundle $Ad(P)$. Let $\bar{\partial}_P$ denote the differential operator of order one defining the holomorphic structure of $Ad(P)$.

Let Ω_M^i denote the holomorphic vector bundle on M given by the holomorphic i -forms.

The holomorphic connection D on P induces a holomorphic connection on $Ad(P)$, which is again denoted by D . In other words,

$$D : Ad(P) \longrightarrow Ad(P) \otimes \Omega_M^1$$

is a first order operator satisfying the Leibniz condition $D(f.s) = \partial f.s + fD(s)$, where f is a smooth function, such that the curvature, $(D + \bar{\partial}_P)^2$, of the connection $D + \bar{\partial}_P$ is a holomorphic section of $\Omega_M^2 \otimes Ad(P)$. This condition implies that the operator D maps a holomorphic sections of $Ad(P)$ to a holomorphic section of $Ad(P) \otimes \Omega_M^1$.

A vector bundle E on M is called *stable* (resp. *semistable*) if for any $\mathcal{O}_M$ -coherent proper subsheaf $0 \neq F \subset E$, with E/F torsion-free, the following condition holds [Ko, Ch. V, §7] :

$$\mu(F) := \deg F / \text{rank} F < \mu(E) := \deg E / \text{rank} E \quad (\text{resp. } \mu(F) \leq \mu(E))$$

A vector bundle is called *quasistable* if it is a direct sum of stable bundles of same μ (slope).

Lemma 2.1. *If $\deg(T/\mathcal{E}_{l-1}) \geq 0$ then the vector bundle $Ad(P)$ on M is semistable. Moreover, if $\deg(T/\mathcal{E}_{l-1}) > 0$, then $Ad(P)$ is quasistable.*

The proof of this lemma is actually contained in [Bi]. However, to be somewhat self-contained, we will give some details of the proof.

Proof of Lemma 2.1. We will first prove that if $\deg(T/\mathcal{E}_{l-1}) \geq 0$ then the bundle $Ad(P)$ is semistable.

Suppose $Ad(P)$ is not semistable. In that case it has a nontrivial Harder-Narasimhan filtration. Let

$$(2.2) \quad 0 = V_0 \subset V_1 \subset V_2 \subset \dots \subset V_{n-1} \subset V_n = Ad(P)$$

be the Harder-Narasimhan filtration of the vector bundle $Ad(P)$.

We restrict the domain of the operator D (which gives the holomorphic connection on $Ad(P)$) to the subsheaf V_1 , and consider the induced operator

$$(2.3) \quad D_1 : V_1 \longrightarrow (Ad(P)/V_1) \otimes \Omega_M^1$$

The Leibniz identity for D implies that $D_1(f.s) = f.D_1(s)$, *i.e.* D_1 is $\mathcal{O}_M$ -linear.

The largest semistable subsheaf (*i.e.* the first nonzero term in the Harder-Narasimhan filtration) of Ω_M^1 is the kernel of the surjective homomorphism

$$q_{l-1} : \Omega_M^1 \longrightarrow \mathcal{E}_{l-1}^*$$

obtained by taking the dual of the inclusion in (1.2). Since the tensor product of two semistable sheaves is again semistable [MR, Remark 6.6 iii], the largest semistable subsheaf of $(Ad(P)/V_1) \otimes \Omega_M^1$ is

$$(2.4) \quad W := (V_2/V_1) \otimes \ker(q_{l-1})$$

The general formula for the degree of a tensor product gives $\mu(W) = \mu(V_2/V_1) + \mu(\ker(q_{l-1}))$. From the property of the Harder-Narasimhan filtration we have that $\mu(V_2/V_1) < \mu(V_1)$. The assumption that $\deg(T/\mathcal{E}_{l-1}) \geq 0$ implies that

$$\mu(\ker(q_{l-1})) = -\mu(T/\mathcal{E}_{l-1}) \leq 0$$

From these we conclude that

$$(2.5) \quad \mu(V_1) > \mu(W)$$

Let E denote the image of D_1 defined in (2.3). Assume that E is not the zero sheaf. Since V_1 is semistable and E is a quotient of V , we have $\mu(E) \geq \mu(V_1)$. On the other hand, since W is the largest semistable subsheaf of $(Ad(P)/V_1) \otimes \Omega_M^1$, we have $\mu(E) \leq \mu(W)$. But this contradicts (2.5). So E must be the zero sheaf, *i.e.* the homomorphism D_1 in (2.3) must be the zero homomorphism. Thus we obtain that the subsheaf V_1 is invariant under the connection D on $Ad(P)$.

Any $\mathcal{O}_M$ coherent sheaf with a holomorphic connection is a locally free $\mathcal{O}_M$ module [B, p. 211, Proposition 1.7]. (This proposition in [B] is stated for integrable connections (D -modules), but the proof uses only the Leibniz rule which is valid for a holomorphic connection.)

Using the Chern-Weil construction of characteristic classes it is easy to show that for a vector bundle E on M , equipped with a holomorphic connection, any Chern class $c_i(E) \in H^{2i}(M, \mathbb{Q})$, $i \geq 1$, vanishes.

So we have $\deg V_1 = 0 = \deg Ad(P)$. If $V_1 \neq Ad(P)$ then $\mu(V_1) > \mu(Ad(P))$. So $Ad(P) = V_1$, *i.e.* the bundle $Ad(P)$ is semistable.

If $Ad(P)$ is stable then obviously it is quasistable. Suppose that $Ad(P)$ is not stable. Then there is a filtration [K, Ch. V, §7, Theorem 7.18]

$$(2.6) \quad 0 = W_0 \subset W_1 \subset W_2 \subset \dots \subset W_{n-1} \subset W_n = Ad(P)$$

such that W_i/W_{i-1} , $1 \leq i \leq n$, is a stable sheaf with $\mu(W_i/W_{i-1}) = \mu(Ad(P))$.

Now, as done in the proof of Proposition 3.4 of [Bi], using the given condition that $\deg(T/\mathcal{E}_{l-1})$ is strictly positive, it is possible to show that the filtration (2.6) splits, *i.e.*

$$(2.7) \quad W_i = W_{i-1} \oplus (W_i/W_{i-1})$$

where W_i/W_{i-1} is a locally free sheaf. The equality (2.7) proves that

$$Ad(P) = \sum_{i=1}^n W_i/W_{i-1}$$

This completes the proof of the lemma. $\square$

In the next section we will use Lemma 2.1 to construct a flat connection on P .

3. EXISTENCE OF A FLAT CONNECTION

Let G be a connected affine algebraic reductive group over $\mathbb{C}$. Let P be a holomorphic principal G bundle over M .

Let $i : U \rightarrow M$ be the inclusion of an open subset such that the complement $X - i(U)$ is an analytic subset of X of codimension at least two. For a $\mathcal{O}_U$ coherent sheaf F on U , the direct image i_*F is a $\mathcal{O}_M$ coherent sheaf. The degree of F is defined to be the degree of i_*F .

We will recall the definition of (semi)stability of P ([RR, Definition 4.7]). Let $U \subset X$ be an open subset with $X - U$ being an analytic set of codimension at least two. Let Q be a parabolic subgroup of G , and let P' be a reduction of the structure group to Q of the restriction of the principal P to the open set U . The principal bundle P is said to be *stable* (resp. *semistable*) if for any such P' , the degree of a line bundle on U associated to P for any character χ on Q dominant with respect to a Borel subgroup contained in Q , is strictly negative (resp. nonpositive).

Let D be a holomorphic connection on P (defined in Section 1). The following theorem is a generalization of Lemma 2.1.

Theorem 3.1. *If $\deg(T/\mathcal{E}_{l-1}) \geq 0$ then P is semistable. Moreover, if $\deg(T/\mathcal{E}_{l-1}) > 0$, then P admits a flat G -connection compatible with the holomorphic structure.*

Proof. Let $P' \subset P$ be a reduction of structure group of P to a maximal parabolic subgroup $Q \subset G$. This reduction is given by a section, σ , of the fiber bundle

$$\rho : P/Q \rightarrow U$$

Let T_{rel} denote the relative tangent bundle for the map ρ .

From Lemma 2.1 of [R] it follows that in order to check that P is semistable it is enough to show that $\deg(\sigma^*T_{\text{rel}}) \geq 0$.

The reduction $P' \subset P$ gives an injective homomorphism $Ad(P') \longrightarrow Ad(P)$ of adjoint bundles on M . The bundle σ^*T_{rel} on M is the quotient bundle $Ad(P)/Ad(P')$.

From Lemma 2.1 we know that if $\deg(T/\mathcal{E}_{l-1}) \geq 0$, the adjoint bundle $Ad(P)$ is semistable. Since G is reductive, the Lie algebra $\mathfrak{g}$ admits a nondegenerate G invariant bilinear form. This implies that $Ad(P) = Ad(P)^*$. Hence $\deg Ad(P) = 0$. Now the semistability of $Ad(P)$ implies that $\deg(Ad(P)/Ad(P')) \geq 0$. This proves that the principal bundle P is semistable.

If $\deg(T/\mathcal{E}_{l-1}) > 0$ then from Lemma 2.1 we know that $Ad(P)$ is quasistable. So from the main theorem of [UY] it follows that the vector bundle $Ad(P)$ admits a Hermitian-Yang-Mills connection. We will denote this Hermitian-Yang-Mills connection by ∇ . This connection ∇ is unique (though the Hermitian-Yang-Mills metric is not unique), and it is irreducible if and only if $Ad(P)$ is stable.

We want to show that this connection ∇ induces a connection on the principal G bundle P .

Let Z_0 denote the connected component of the center of G containing the identity element (the center has finitely many components). Define

$$G_0 = G/Z_0$$

which is a semisimple group. The group G_0 acts on the Lie algebra $\mathfrak{g}$ (of G) by conjugation and gives an homomorphism

$$(3.2) \quad \theta : G_0 \longrightarrow GL(\mathfrak{g})$$

which has a finite group as the kernel.

From a theorem of Chevalley [H, Theorem 11.2] we know that there is a linear representation of the group $GL(\mathfrak{g})$

$$(3.3) \quad \phi : GL(\mathfrak{g}) \longrightarrow GL(V)$$

in a vector space V over $\mathbb{C}$ and a line L in V such that

$$\phi \circ \theta(G_0) = \{g \in GL(\mathfrak{g}) \mid \phi(g)(L) = L\}$$

Since G_0 is semisimple, it does not have any nontrivial character. This implies that $\phi \circ \theta(G_0)$ fixes the line L point-wise. Let $0 \neq v \in L$ be a nonzero vector. So the isotropy subgroup of v for the action of $GL(\mathfrak{g})$ on V is precisely $\phi \circ \theta(G_0)$. It is not difficult to see that the homomorphism ϕ can be so chosen that it maps the center of the Lie algebra of $GL(\mathfrak{g})$ into the center of the Lie algebra of $GL(V)$. We will choose ϕ such that it satisfies this condition.

Let $q : G \longrightarrow G_0$ denote the obvious projection. Let $P(G_0)$ denote the principal G_0 bundle on M obtained by extending the structure group of P to G_0 using the homomorphism q . The vector bundle $Ad(P)$, which can be identified with a principal $GL(\mathfrak{g})$ bundle, is obtained by extending the structure group of $P(G_0)$ to $GL(\mathfrak{g})$ using the homomorphism θ defined in (3.2).

Using the homomorphism ϕ we may extend the structure group of $Ad(P)$ to $GL(V)$. The vector bundle on M associated to this principal $GL(V)$ bundle for the natural action of $GL(V)$ on the vector space V will be denoted by E .

The Hermitian-Yang-Mills metric on the vector bundle $Ad(P)$ gives a reduction of the structure group of $Ad(P)$ to $U(\mathfrak{g})$, a maximal compact subgroup of $GL(\mathfrak{g})$. Since the image $\phi(U(\mathfrak{g}))$ is a compact subgroup of $GL(V)$, it is contained in some maximal compact subgroup of $GL(V)$. So the reduction of $Ad(P)$ to $U(\mathfrak{g})$ will induce a reduction of E to a maximal compact subgroup of $GL(V)$ (*i.e.* the vector bundle E will be equipped with a hermitian metric) such that the connection on E obtained by extending the connection ∇ (on $P(\mathfrak{g})$) is the hermitian connection (for the hermitian metric on E). Since the metric on $Ad(P)$ is a Hermitian-Yang-Mills metric, the metric on E obtained above is also a Hermitian-Yang-Mills metric. Indeed, the Hermitian-Yang-Mills condition of the connection on $Ad(P)$ implies that the curvature is a 2-form on M with values in the center of the endomorphism bundle $Ad(P)^* \otimes Ad(P)$. Since ϕ maps the center of the Lie algebra of $GL(\mathfrak{g})$ into the center of the Lie algebra of $GL(V)$, the induced connection on E is a Hermitian-Yang-Mills connection. Let ∇' denote the Hermitian-Yang-Mills connection on E obtained this way.

Since $\theta(G_0)$ fixes the vector v , this vector v will give a nowhere zero section of E (since E is obtained by extending the structure group of P to $GL(V)$); let s denote this section of E .

The holomorphic connection D on P will induce a holomorphic connection on E . As we noted in Section 2, using the Chern Weil construction it is easy to see that the existence of a holomorphic connection on E implies that any Chern class, $c_i(E)$, $i \geq 1$, vanishes.

From [Ko, Ch. IV, Corollary 4.13] it follows that Hermitian-Yang-Mills connection ∇' is a flat connection. Since ∇' is a flat unitary connection, any holomorphic section of E must be a flat section for the connection ∇' . Indeed, the Laplacian of a flat unitary connection operator is twice the Laplacian of the Dolbeault operator. In particular, the space of harmonic sections for these two Laplacians coincide. Since any holomorphic section is a harmonic section for the Dolbeault Laplacian, it must be a flat section. So, in particular, the section s is flat.

Let $P(\mathfrak{g})$ denote the principal $GL(\mathfrak{g})$ obtained by extending the structure group of P to $GL(\mathfrak{g})$ using the homomorphism $\theta \circ q$. Since $P(\mathfrak{g})$ is an extension, the fiber bundle, $P(\mathfrak{g})/G$, with fiber $GL(\mathfrak{g})/G$ has a natural section (which gives the reduction of the structure group of $P(\mathfrak{g})$ to G). Let α denote this section of $P(\mathfrak{g})/G$.

Since $\theta(G_0)$ fixes v for the homomorphism ϕ in (3.3), we have an embedding of the fiber bundle $P(\mathfrak{g})/G$ in the total space of E (given by the orbit of v for the action of $GL(\mathfrak{g})$ using ϕ).

It is easy to see that the image of the section α by the above embedding of $P(\mathfrak{g})/G$ in E is precisely the section s .

Now, since s is a flat section for the connection ∇' , the connection ∇ induces a G_0 connection on the principal G_0 bundle $P(G_0)$ on M as follows : Let

$$p : P(G_0) \longrightarrow P(\mathfrak{g})$$

denote the holomorphic map induced by θ in (3.2). Take $x \in p(P(G_0))$, and let $v \in T_x P(\mathfrak{g})$ be a horizontal vector for the connection ∇ on $P(\mathfrak{g})$. Let w be the image

of v by the differential of the map

$$P(\mathfrak{g}) \longrightarrow E$$

induced by the homomorphism ϕ in (3.3). Since s is flat, the tangent vector w lies in the submanifold of the total space of E given by the section s . But this implies that the tangent vector v lies in the image of $TP(G_0)$ under the map given by the differential of p . Thus the connection ∇ on $P(\mathfrak{g})$ induces a connection on the principal bundle $P(G_0)$ with structure group G_0 . We will call this connection on $P(G_0)$ as ∇_0 . Since ∇ is a flat connection, ∇_0 is also flat.

The commutator subgroup $G' := [G, G] \subset G$ is a semisimple group, and the restriction of q to G' is a surjective homomorphism with a finite kernel. So their Lie algebras are isomorphic. So

$$(3.4) \quad G = Z_0.G'$$

with a finite intersection $\Gamma := Z_0 \cap G'$.

The abelian Lie group Z_0/Γ is a product of copies of $\mathbb{C}^*$, since G is assumed to be affine. Let

$$f : G \longrightarrow Z_0/\Gamma$$

denote the obvious projection (obtained from (3.4)).

Let $P(f)$ denote the principal Z_0/Γ bundle on M obtained by extending the structure group of P using the homomorphism f .

The holomorphic connection D on P induces a holomorphic connection on $P(f)$, which we will denote by $D(f)$.

Any holomorphic line bundle on M admitting a holomorphic connection actually admits a compatible flat connection. Indeed, if ∂ is a holomorphic connection on a holomorphic line bundle L whose holomorphic structure is given by the operator $\bar{\partial}$, then the curvature $(\partial + \bar{\partial})^2$ is a holomorphic 2-form which is exact (since the cohomology class represented by it is of the type $(1, 1)$). So it is of the form $\partial\beta$, where β is a $(1, 0)$ -form. The new connection

$$\partial - \beta + \bar{\partial}$$

on L is a flat connection compatible with the holomorphic structure.

Recall that Z_0/Γ is a product of copies of $\mathbb{C}^*$. In view of the above remark, the existence of the holomorphic connection $D(f)$ implies that the principal Z_0/Γ bundle $P(f)$ admits a flat connection. Let ∇_1 be a flat connection on $P(f)$.

Since the exact sequence of the Lie algebras

$$0 \longrightarrow \mathfrak{z}_0 \longrightarrow \mathfrak{g} \longrightarrow \mathfrak{g}_0 \longrightarrow 0$$

has a natural splitting (given by the Lie algebra of G'), the two flat connections ∇_0 and ∇_1 combine together to induce a flat G connection on P as follows : The horizontal subspace of the tangent space at a point $p \in P$ is defined to be the intersection of the inverse images of the horizontal subspaces of $P(G_0)$ and $P(f)$ (horizontal subspaces for the flat connections ∇_0 and ∇_1 respectively) for the obvious projections of P onto $P(G_0)$ and $P(f)$ respectively. The integrability of ∇_0 and ∇_1 will imply that the

connection on P obtained above is actually flat. This completes the proof of the theorem. $\square$

REFERENCES

- [B] A. Borel et al. : Algebraic D -modules. Perspectives in Mathematics, Vol. 2 (Ed. J. Coates, S. Helgason), Academic Press, 1987.
- [Bi] I. Biswas : On Harder-Narasimhan filtration of the tangent bundle. To appear in Comm. Anal. Geom.
- [H] J. Humphreys : Linear algebraic groups. Graduate Texts in Math. 21, Springer-Verlag, New York Heidelberg Berlin, 1975.
- [Ko] S. Kobayashi : Differential geometry of complex vector bundles. Publications of Math. Soc. of Japan, Iwanami Schoten Pub. and Princeton University Press, 1987.
- [M1] S. Murakami : Sur certains espaces fibrés principaux admettant des connexions holomorphes. Osaka Math. Jour. **11** (1959) 43-62.
- [M2] S. Murakami : Sur certains espaces fibrés principaux holomorphes dont le groupe est abélien connexe. Osaka Math. Jour. **13** (1961) 143-167.
- [M3] S. Murakami : Harmonic connections and their applications. SEA Bull. Math. Special Issue (1993) 101-103. World Sci. Pub. Company.
- [MR] V. Mehta, A. Ramanathan : Semistable sheaves on projective varieties and their restriction to curves. Math. Ann. **258** (1982) 213-224.
- [RR] S. Ramanan, A. Ramanathan : Some remarks on the instability flag. Tôhoku Math. J. **36** (1984) 269-291.
- [R] A. Ramanathan : Stable principal bundles on a compact Riemann surface. Math. Ann. **213** (1975) 129-152.
- [UY] K. K. Uhlenbeck, S. T. Yau : On the existence of Hermitian-Yang-Mills connections in stable vector bundles. Comm. Pure Appl. Math. **39** (1986) 257-293.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400005, INDIA

Current address: Université de Grenoble 1, Institut Fourier, 100 rue des Maths, B.P. 74, 38402 Saint-Martin-d'Hères, FRANCE

E-mail address: `indranil@math.tifr.res.in` and `biswas@puccini.ujf-grenoble.fr`